

π, K, p and $\bar{p}$ production from Au+Au collisions at $\sqrt{s_{NN}} = 62.4$ GeV

Lijuan Ruan^{†‡} (for the STAR Collaboration[§])

Dept. of Modern Physics, University of Science and Technology of China, Hefei, Anhui, China, 230026

Abstract. The preliminary results of $\pi^\pm, K^\pm, p$ and $\bar{p}$ spectra are reported from Au+Au collisions at $\sqrt{s_{NN}} = 62.4$ GeV. Particle identification is from the Time Projection Chamber and Time-of-Flight system at STAR. The nuclear modification factor R_{CP} for mesons ($\pi^\pm, K^\pm$) and baryons ($p, \bar{p}$) will also be discussed.

In central 200 GeV Au+Au collisions at RHIC, hadron production at high p_T ($p_T > 5$ GeV/c) has been observed to be suppressed relative to that from p+p collisions scaled by the number of binary collisions and the measured nuclear modification factor is significantly below unity [1, 2]. This suppression has been interpreted as energy loss of the energetic partons traversing the produced hot and dense medium [3]. At intermediate p_T ($2 < p_T < 5$ GeV/c), the degree of suppression depends on particle species. The nuclear modification factors of baryons (protons and lambdas) are significantly higher than those of mesons (pions, kaons) [4, 5], which can be explained by coalescence of constituent quarks at hadronization from a collective partonic system [6]. In Au+Au collisions at 62 GeV, the bulk properties created was expected to be different from those at 200 GeV. The smaller gluon density was expected [7]. In such an environment, we'd like to study the energy loss behavior and the particle composition at intermediate and high p_T . In this paper, the particle spectra and their nuclear modification factors of π, K, p and $\bar{p}$ will be reported. Also presented are the ratios of protons over pions.

1. Analysis methods and results

At STAR, charged particles can be identified up to $p_T \sim 1.1$ GeV/c by measuring their ionization energy loss (dE/dx) in the Time Projection Chamber (TPC) [10]. The dE/dx resolution was calibrated to be better than 8% in 62.4 GeV Au+Au collisions for tracks with 70 cm in length inside the TPC. At momentum $p > 3$ GeV/c, the dE/dx of $\pi^\pm$ has a $\sim 2\sigma$ separation from those of $K^\pm$ and $p(\bar{p})$ due to the relativistic rise of pion dE/dx . Thus charged pions can be identified at $3 < p_T < 10$ GeV/c [9]. The dE/dx measurement in Fig. 1 (left) uses a normalized dE/dx : $n\sigma_X^Y = \ln((dE/dx)^Y/I_{0X})/\sigma_X$, where X, Y can be $e^\pm, \pi^\pm, K^\pm$ or $p(\bar{p})$, $(dE/dx)^Y$ is the measured dE/dx of particle Y , I_{0X}

[†] This work is supported in part by NSFC (10155002).

[‡] rlj@mail.ustc.edu.cn

[§] For the full author list and acknowledgements see Appendix “Collaborations” in this volume.




Figure 1. (left) dE/dx distribution normalized by pion dE/dx and offset by ± 5 for positive and negative charged particles at $4 < p_T < 4.5$ GeV/c, respectively. The distribution is from minimum-bias Au+Au collisions at 62.4 GeV [9]. (right) m^2 versus $n\sigma_\pi$ at $2 < p_T < 4$ GeV/c in minimum-bias Au+Au collisions at 62.4 GeV.

is the expected dE/dx of particle X , and σ_X is the dE/dx resolution of particle X . With a perfect calibration, the $n\sigma_\pi^\pi$ distribution should be a normal Gaussian while $n\sigma_\pi^p, n\sigma_\pi^K$ should be a Gaussian peaked at negative value. The method was introduced in [9]. A prototype multi-gap resistive plate chamber time-of-flight system (TOFr) [10] has been installed since 2003 with the coverage $-1 < \eta < 0$ in pseudorapidity. With intrinsic timing resolution 85 ps [11], it extends particle identification up to $p_T \sim 3$ GeV/c for p and $\bar{p}$ and 1.6 GeV/c for $\pi^\pm$ and $K^\pm$. By the combination of $m^2 = p^2(1/\beta^2 - 1)$ from TOFr and $n\sigma_\pi, n\sigma_p$ information from TPC, $\pi^\pm, p$ and $\bar{p}$ can be identified up to $p_T \sim 5$ GeV/c and $K^\pm$ can be identified to at least 3 GeV/c, where β is the velocity. This method has been introduced in [12]. Fig. 1 (right) shows m^2 versus $n\sigma_\pi$.

A total of 6.8 million events after vertex cut $|Vz| < 30$ cm were used for the analysis, where Vz is the z position of the vertex. Centrality tagging of Au+Au collisions was based on the charged particle multiplicity in $-0.5 < \eta < 0.5$, measured by the TPC. The minimum-bias (0-80%) events were divided into four centralities: most central 10%, 10 – 20%, 20 – 40% and 40 – 80% of the hadronic cross section.

From TOFr, the raw yields of $\pi^\pm, K^\pm, p$ and $\bar{p}$ are obtained from Gaussian fits to the distributions in $m^2 = p^2(1/\beta^2 - 1)$ in each p_T bin w/o $n\sigma$ cut. Acceptance and efficiency were studied by Monte Carlo simulations and by matching TPC track and TOFr hits in real data. From TPC, the raw yields of $\pi^\pm$ were extracted from dE/dx distribution. The efficiency due to the additional $n\sigma$ cut was also taken into account. Weak-decay feeddown (e.g. $K_s^0 \rightarrow \pi^+\pi^-$) to pions was not corrected for, which was estimated to contribute $\sim 12\%$ at $p_T < 1$ GeV/c and $\sim 5\%$ at higher p_T to pion yields [11]. Inclusive p and $\bar{p}$ production is presented without hyperon feeddown correction either. p and $\bar{p}$ from hyperon decays have the same detection efficiency as primary p and $\bar{p}$ [13, 14] and the contributions to the inclusive p and $\bar{p}$ yield range from $\sim 20\%$ to $\sim 40\%$ from p+p, d+Au to Au+Au collisions [11, 13, 14].

The invariant yields $d^2N/2\pi p_T dp_T dy$ of π^-, K^- and $\bar{p}$ at mid-rapidity, after the efficiency correction, are shown as symbols in Fig. 2 (left) for 62.4 GeV minimum bias Au+Au collisions. In the overlapping p_T region, the results from TOFr and from TPC are consistent. Fig. 2 (middle) shows the anti-particle to particle ratios versus p_T .




Figure 2. (left) The spectra of π^- , K^- and $\bar{p}$ versus p_T . (middle) π^-/π^+ , K^-/K^+ and $\bar{p}/p$ ratios versus p_T . (right) The π^- rapidity density versus N_{part} .

Within the errors, π^-/π^+ , K^-/K^+ and $\bar{p}/p$ are flat with p_T . The average anti-particle to particle ratios were obtained with a fit of constant value: $\pi^-/\pi^+ = 1.02 \pm 0.01$, $K^-/K^+ = 0.84 \pm 0.01$ and $\bar{p}/p = 0.46 \pm 0.01$. The errors are statistical. The systematical errors are similar to those presented at [11, 13]. At 200 GeV, $\bar{p}/p = 0.77 \pm 0.05$. The decrease of $\bar{p}/p$ ratio at 62 GeV indicates the increase of the baryon chemical potential. The mid-rapidity yield dN/dy of π^- was extracted with a Bose-Einstein fit [13]. Fig. 2 (right) shows the dN/dy of π^- versus N_{part} in Au+Au collisions at 62 and 200 GeV [13]. The dN/dy at 62 GeV is a factor of ~ 1.5 smaller than that at 200 GeV [13].




Figure 3. (left) R_{AA} of $\pi^\pm$ versus p_T in Au+Au collisions at 62.4 GeV. Also shown are the two curves from theoretical predictions for 62.4 GeV [7]. Reference spectrum of p+p is from ISR [8]. (middle) R_{CP} of $\pi^\pm$ and h . (right) R_{CP} of $\pi^\pm$, $K^\pm$ and $p + \bar{p}$.

Nuclear effects on hadron production in Au+Au collisions were measured through comparison to the p+p spectrum scaled by the number of underlying nucleon-nucleon inelastic collisions (N_{bin}). Fig. 3 (left) shows the R_{AA} versus p_T from our measurement in most central 0-10% Au+Au collisions at 62.4 GeV. Also shown in this plot is the R_{AA} in most central 0-10% Au+Au collisions at 200 GeV from PHENIX measurement [2]. It's evident that the suppression exists at 62.4 GeV and that the magnitude of suppression is smaller than that at 200 GeV. Fig. 3 (middle) shows R_{CP} of charged hadron (h) and $\pi^\pm$ from 62.4 GeV Au+Au collisions. R_{CP} of h [15] is larger than that of $\pi^\pm$ at $2 < p_T < 5$ GeV/c and approaches that of $\pi^\pm$ at $5 < p_T < 7$ GeV/c. This may indicate the disappearance of particle-species dependence of nuclear modification factor at high p_T . Fig. 3 (right) shows R_{CP} of $\pi^\pm$, $K^\pm$ and $p + \bar{p}$ from 62.4 GeV Au+Au collisions. R_{CP} of protons seems to follow N_{bin} scaling at intermediate p_T and be larger than those of pions. The statistic for kaons is too poor to address physics issues.

Fig. 4 shows the ratios of p/π^+ and $\bar{p}/\pi^-$ as a function of p_T in Au+Au collision at

62.4 GeV. The p/π^+ and $\bar{p}/\pi^-$ ratios are observed to be a factor of $2 \sim 3$ higher than those in p+p collisions at similar energy at $2 < p_T < 4$ GeV/c. p/π^+ and $\bar{p}/\pi^-$ ratios reach maximum at around 2 GeV/c and seem to decrease with increasing p_T at higher p_T . This is consistent with the trends of R_{CP} of h and π . At $p_T \sim 5$ GeV/c, p/π^+ and $\bar{p}/\pi^-$ ratios in 62 GeV Au+Au collisions are close to those in p+p collisions [16]. This may indicate that fragmentation mechanism dominates above this p_T region.




Figure 4. The ratios of p/π^+ (left) and $\bar{p}/\pi^-$ (right) as a function of p_T in Au+Au collision at 62.4 GeV. Also shown are the ratios in p+p collisions at 53 GeV [16].

2. Summary

We report the STAR preliminary results of $\pi^\pm, K^\pm, p$ and $\bar{p}$ spectra from 62.4 GeV Au+Au collisions. $\pi^\pm$ and $p(\bar{p})$ were identified up to $p_T \sim 7$ GeV/c and 5 GeV/c, respectively. At this beam energy, $\bar{p}/p = 0.46 \pm 0.01$. A significant suppression for pions is observed for the most central collisions, but the effect is weaker than that observed in 200 GeV central Au+Au collisions [2] at $p_T < 7$ GeV/c. At intermediate p_T , the nuclear modification factor R_{CP} of h is 20% higher than that of π , R_{CP} of $p + \bar{p}$ is larger than that of π , and the ratios of p/π^+ and $\bar{p}/\pi^-$ are a factor of $2 \sim 3$ higher than those in p+p collisions at similar energies [16].

References

- [1] STAR Collaboration, J. Adams *et al.*, Phys. Rev. Lett. **91**, 172302 (2003).
- [2] PHENIX Collaboration, S.S. Adler *et al.*, Phys. Rev. Lett. **91**, 072301 (2003); PHENIX Collaboration, S.S. Adler *et al.*, Phys. Rev. Lett. **91**, 241803 (2003).
- [3] M. Gyulassy *et al.*, nucl-th/0302077.
- [4] STAR Collaboration, J. Adams *et al.*, Phys. Rev. Lett. **92**, 052302 (2004).
- [5] PHENIX Collaboration, K. Adcox *et al.*, Phys. Lett. B **561**, 82 (2003); PHENIX Collaboration, S.S. Adler *et al.*, Phys. Rev. Lett. **91**, 172301 (2003).
- [6] R.C. Hwa *et al.*, Phys. Rev. C **67**, 034902 (2003); R.J. Fries *et al.*, Phys. Rev. C **68**, 044902 (2003); V. Greco *et al.*, Phys. Rev. Lett. **90**, 202302 (2003).
- [7] I. Vitev, nucl-th/0404052, curve shown is calculation with $dN_g/dy=650$; X.N. Wang, Phys. Rev. C **70**, 031901 (2004).
- [8] PHENIX Collaboration, D. d'Enterria *et al.*, The Proceedings in Hot Quarks 2004.
- [9] STAR Collaboration, Z. Xu *et al.*, The Proceedings in DPF 2004, nucl-ex/0411001.
- [10] M. Anderson *et al.*, Nucl. Instr. Meth. A **499**, 659 (2003); B. Bonner *et al.*, Nucl. Instr. Meth. A **508**, 181 (2003); M. Shao *et al.*, Nucl. Instr. Meth. A **492**, 344 (2002).
- [11] STAR Collaboration, J. Adams *et al.*, nucl-ex/0309012.
- [12] STAR Collaboration, M. Shao *et al.*, The Proceedings in Hot Quarks 2004, nucl-ex/0411035.
- [13] STAR Collaboration, J. Adams *et al.*, Phys. Rev. Lett. **92**, 112301 (2004).
- [14] STAR Collaboration, C. Adler *et al.*, Phys. Rev. Lett. **87**, 262302 (2001).
- [15] STAR Collaboration, J. Dunlop *et al.*, RHIC & AGS Annual Users Meeting, May 10-14, 2004.

[16] B. Alper *et al.*, Nucl. Phys. B **100**, 237 (1975).